

Figure 1a. LCMS analysis of recombinant peptide variants

(A) MM-416776

(B) MD-915







Figure 1b: LCMS analysis of synthetic MD-1100 (Total Ion Chromatograph (TIC))




**Figure 1c: LCMS analysis (Total Ion Chromatograph of
blank used in MD-1100 analysis)**



**Figure 2. Chemically synthesized peptides in the
 Intestinal GC-C Receptor Activity Assay**



**Figure 3a. MM-416776 vs Zelnorm® in an acute Mouse
Gastrointestinal Transit Model (GIT)**



**Figure 3b: MD-1100 vs. Zelnorm® in an acute Mouse
 Gastrointestinal Transit Model**





Figure 4a. Purified MD-915 and MM-416776 in GIT Model




Figure 4b. Chemically Synthesized Peptides in GIT Model

Figure 4c. Chronic vs. Acute Dosing in GIT Assay



**Figure 5a. MM-416776 vs Zelnorm® in a Mouse Intestinal
Secretion Model**



**Figure 5b: MD-1100 vs Zelnorm® in Mouse Intestinal
Secretion Model**







Figure 6a. Recombinantly generated MD-915 and MM-416776 in Mouse Intestinal Secretion Model



**Figure 6b. Chemically synthesized peptides in Mouse
 Intestinal Secretion Model**



**Figure 7: Effect of MD-1100 on pain in a rat TNBS
 Colorectal Distention Assay**



* $p < 0.05$ as compared to "vehicle" value

**Figure 8a: Visceral Antinociceptive Effects of MD-915 in a
 Mouse Writhing Assay**




Figure 8b: Visceral Antinociceptive Effects of MD-1100 in a Mouse Writhing Assay






Figure 9: Competitive Radioligand Binding of MD-1100




**Figure 10a: Minimum Systemic Absorption of MD-1100
 (based on ELISA)**



* Limit of detection 0.061 µg/ml (40 nM)
 Dosing at 10 mg/kg

**Figure 10b: Minimum Systemic Absorption of MD-1100
 (based on LC/MS)**



- Limit of detection 0.00063 µg/mL (0.6 nM)
- Dosing at 10 mg/kg